

## GUIDELINE FOR VAGINAL BIRTH AFTER ONE PREVIOUS LOW SEGMENT CESAREAN SECTION

The College of Midwives of British Columbia supports registered midwives in providing primary care for women planning a vaginal birth after one previous low-segment cesarean section (VBAC).<sup>1</sup>

The rate of successful VBAC in Canada is between 50-85% (SOGC, 2005). In the 2009 -10 BC Perinatal Database Registry, BC midwives had a VBAC success rate of 80%.<sup>2</sup>

### **Risk of Uterine Rupture**

Current evidence supports women in choosing vaginal birth after cesarean section, despite a somewhat increased risk of uterine rupture, a complication with potentially serious consequences for mother and newborn.

Previous cesarean section is a factor in somewhat less than half of all reported cases of uterine rupture. Previous vaginal birth or previous labour before a primary cesarean delivery can decrease the risk of uterine rupture in a subsequent trial of labour (Marshak et al, 2000, Algert et al, 2008). Varying rates of uterine dehiscence<sup>3</sup> and rupture<sup>4</sup> in the presence of a previous cesarean section scar are reported in the literature. One difficulty in looking at the literature is that the rate of rupture for VBACs with induced or augmented labour is often higher than for spontaneous labour, but these rates may be combined and reported as an overall rate. Whether the previous cesarean incision had a locked or unlocked single layer closure or double layer closure, another factor thought to affect the rate of rupture, is also not always addressed. Another difficulty is that the rate of catastrophic rupture, where the life of mother and infant are in serious jeopardy, is more difficult to determine as this event is often included with the more common and much less worrisome dehiscence. Overall VBAC research remains of limited value in predicting risk of symptomatic uterine rupture.

The general consensus on the rate of uterine rupture in spontaneous labour with one lower segment incision is 0.4%, and in augmented labour is 0.9%. The overall rate of uterine rupture reported in the peer reviewed VBAC literature is 0.7%. The rate for women with 2 or more cesarean scars increases to 0.9% (Landon, 2006). Overall rates of uterine rupture for VBAC have been reported as low as 0.2% to as high as 0.7% depending on which factors in a woman's health history were included in studies (Nahum, 2012).

---

<sup>1</sup> The College's *Indications for Discussion, Consultation and Transfer of Care* requires the plan of care for a VBAC client with a history of one prior lower segment incision be discussed with another midwife. A physician consultation is required for a client with a history of more than one previous cesarean section..

<sup>2</sup> BC Perinatal Database Registry Annual Report March 31, 2010.

<sup>3</sup> Scar dehiscence is the breakdown and reopening of the old cesarean scar. Most dehiscences involve minor tearing around the scar, are asymptomatic and heal well. Many go undetected.

<sup>4</sup> A true uterine rupture in a VBAC is a scar dehiscence that is large enough to need surgical repair. It is almost always symptomatic, with the most common first indicator being fetal distress. Maternal shock from blood loss is also possible.

The midwife should inform a client wishing to have a VBAC in either a hospital or out-of-hospital setting, of the risk of uterine rupture. The informed choice process for considering VBAC, including choice of birth place, or elective repeat cesarean delivery should be evidence-based, should minimize bias and include the preferences of pregnant women. A copy of this guideline or an alternate written client handout should be offered.

### **Considerations for Choice of Birth Place**

Many women with a history of previous cesarean section with no contraindications to VBAC will be comfortable having a subsequent vaginal birth in hospital with midwifery care. Hospital birth in a facility with cesarean section capability will likely offer the most timely access to emergency cesarean section in the event of uterine rupture. However, some VBAC women will come to midwives requesting home birth. Clients with the following conditions may be candidates for vaginal birth in hospital, but should be advised that they are NOT suitable candidates for a home birth:

- ❖ History of cesarean section at or before 26 weeks
- ❖ History of single layer closure (Bujold et al, 2010, Zelop, 2012)<sup>5</sup>
- ❖ History of infection or impaired uterine scar healing
- ❖ Inter-delivery interval of 18-24 months or less (Bujold et al, 2010)<sup>6</sup>
- ❖ Ballotable head in active labour in current pregnancy
- ❖ Prolonged active phase of labour in current pregnancy

Midwives should always request and review operative reports of previous cesarean sections as well as records of previous obstetric history.

### **Time and Distance – An additional risk**

Despite the relatively small risk, true uterine rupture is a major obstetrical complication with potentially grave consequences for both mother and newborn. Being able to access a cesarean section quickly is very important. There is evidence that suggests that if birth cannot occur between 10 to 37 minutes after a true uterine rupture then fetal death or serious permanent injury is a likely outcome (Nahum, 2012). When planning a VBAC at home, the time it will take to travel to a hospital with cesarean section capabilities must be considered in the light of this small window of time in which one must initiate a cesarean when there is a uterine rupture. Distance to a hospital with cesarean section capability and road and weather conditions are all factors that need to be considered by any woman thinking about planning a VBAC either in an out-of-hospital setting or in a hospital without cesarean section capabilities.

Midwives should discuss these additional risks with their clients and advise them about their local hospital's ability to respond to emergency situations.

### **Client Informed Choice Discussions**

Antenatal client informed choice discussions and teaching should include:

- a) concerns related to previous cesarean section;
- b) possibility of repeat cesarean section;
- c) signs and symptoms of uterine rupture;

---

<sup>5</sup> Locked single layer closure may be associated with greater risk of uterine rupture than an unlocked single layer closure or a double layer closure. However the evidence is limited and further research is required. A two layer uterine closure is recommended for all women considering a future trial of labour after a previous cesarean delivery.

<sup>6</sup> VBAC candidates with inter-delivery interval of 18-24 months should be advised that they may be at increased risk of uterine rupture however the evidence on this time frame remains unclear.

- d) reasons to consult or transport to hospital during labour;
- e) distance to a hospital with cesarean section capabilities;
- f) community standards for vaginal birth after cesarean section<sup>7</sup>.

Documentation of the discussion between the midwife and client of the risks and benefits of VBAC should be made on the antenatal record.

### **Labour Management**

Labour management for VBAC should include:

- a) regular assessment of labour progress and maternal health, with particular awareness of the signs of impending uterine rupture;
- b) regular assessment of fetal health according to the College's *Guidelines for Fetal Health Surveillance in Labour*;
- c) reasonable progress in effacement, dilation and descent every 2-4 hours in active labour;
- d) if labour is occurring out-of-hospital, initiation of transport to hospital if
  - there are concerns about maternal or fetal well-being, including any signs which could indicate impending or actual uterine rupture,
  - the first stage of labour is prolonged, or
  - there is minimal progress in the first hour of active second stage pushing or within two hours of full dilation; and
- e) close observation of blood loss in the hour immediately following delivery of the placenta.

### **Signs that may occur with Impending Uterine Rupture**

- ❖ Inadequate progress (of cervical dilation or descent) despite good contractions
- ❖ Incoordinate uterine activity
- ❖ Restlessness and anxiety
- ❖ Lower abdominal pain or suprapubic tenderness between contractions
- ❖ Hematuria
- ❖ Bandl's ring

### **Signs that may occur with Complete or Partial Rupture**

Midwives must be aware of the signs and symptoms that may indicate uterine rupture in labour. (Rupture of the uterus prior to labour is a rare event and usually involves a classical scar rather than a low-segment scar.)

- ❖ Sudden fetal distress-tachycardia, bradycardia or decelerations (prolonged, late or recurrent variable decelerations are often the first signs of uterine rupture)
- ❖ Unusual abdominal/uterine pain
- ❖ Cessation of contractions or incoordinate uterine activity
- ❖ Unexplained vaginal bleeding
- ❖ A sudden onset of maternal tachycardia and hypotension
- ❖ Excessive fetal movement
- ❖ Fetal parts may be easily palpated through the abdominal wall
- ❖ Presenting part may be higher than previously palpated
- ❖ Signs or symptoms of shock

---

<sup>7</sup> As informed by such documents as SOGC guidelines.

Signs or symptoms of impending or actual uterine rupture in a VBAC client are indications for immediate transport to hospital and physician consultation. Transfer of care will be required unless rupture is ruled out on consultation. If uterine rupture is suspected, the midwife initiating transport or seeking consultation should ask the hospital to prepare for an emergency cesarean section.

Incomplete rupture may also cause postpartum hemorrhage following vaginal birth. If blood loss and/or signs of shock are unexplained in the immediate postpartum, or the mother fails to respond to treatment, the possibility of rupture should be considered.

## References

Algert, CS, Morris, JM, Simpson, JM, Ford, JB, Roberts, CL. Labor before a primary caesarean delivery: reduced risk of uterine rupture in a subsequent trial of labor for vaginal birth after caesarean. *Obstetrics and Gynecology* 2008; 112(5):1061-6.

Appleton, B., Targett, C., Rasmussen, M., Readman, E., Sale, F., and Permezel, M., Vaginal birth after cesarean section: an Australian multicentre study. VBAC Study Group, Feb 2000;40(1) *Aust N Obstet Gynecol*, 933-6.

Blumenfeld, YJ, Caughey, AB., El- Sayed YY, et al. Single versus double-layer hysterotomy closure at primary caesarean delivery and bladder adhesions. *BJOG*,2010;117:690.

Bujold, E., Bujold, C., Hamilton, E.F., Harel, F., Gauthier, R.J. The impact of single-layer or double-layer closure on uterine rupture, *Am J Obstet Gynecol*, 2002;186:1326-30.

Bujold, E., Goyet, M., Marcoux, S, et al. The Role of Uterine Closure in the Risk of Uterine Rupture. *Obstet Gynecol*. 2010;116(1):43-50.

Bujold E et al, Risk of uterine rupture associated with an interdelivery interval between 18 and 24 months. *Int J Obstet Gynecol*. 2010;115(5):1003-6.

CAESAR study collaborative group. Caesarean section surgical techniques; a randomized factorial trial (CAESAR). *GJOG*, 2010; 117:1366.

Enkin M., Keirse M, Neilson J., Crowther C., Duley L., Hodnett E., & Hofmeyr J., 3<sup>rd</sup> Ed A Guide to Effective Care on Pregnancy and Childbirth. (2000) Oxford University Press: Oxford UK

Landon MB, Spong CY, Thom E, et al. Risk of uterine rupture with a trial of labor in women with multiple and single prior cesarean delivery. *Obstet Gynecol*. 2006;108(1):12-20.

Landon MB,, Hauth JC,, Leveno KJ. et al. Maternal and Perinatal Outcomes Associated with a Trial of Labor after Prior Cesarean Delivery. *N Eng J Med*. 2004; Vol. 351 No. 25.

Lydon-Rochelle, M., Holt, V.L., Easterling, T.r., Martin, D.P. Risk of uterine rupture during labour among women with a prior cesarean delivery, *N Engl J Med*, 2001;345:3-8.

Marshak, J.D., Cooperman, B.S., Fried, W.B. and Shi, Q.. (), Prognostic indicators for successful vaginal birth after cesarean delivery, *Obstet Gynecol*.. 2000; S38:94 4(2)

Nahum, G., MD, FACOG, FACS; Chief Editor: Chelmow, D., MD. Uterine Rupture in Pregnancy. *Medscape Updated*: Jul 13, 2012. Accessed July 31, 2012

National Institutes of Health (NIH), Consensus Development Conference on Vaginal Birth After Cesarean: New Insights. Final Panel Statement, *Obstet Gynecol ("The Green Journal")*-2010; Vol. 115, No. 6, June 2010, pp. 1279-1295.

Rizwan N; Abbasi RM; Uddin SF Uterine rupture, frequency of cases and fetomaternal outcome. *J Pak Med Assoc.* 2011; 61(4):322-4

Roberge S et al. Single- versus double-layer closure of the hysterotomy incision during cesarean delivery and risk of uterine rupture. *Int J Obstet Gynecol.*, 2011;115(1):5-10

Schipp, T., Zelop, C.M. Repke, J.T., Cohen, A., and Lieberman, E. Interdelivery interval and risk of symptomatic uterine rupture, *Obstet Gynecol*, 2001;97(2) 175 -7.

Shimonovitz, S., Botosneano, A., Hochner-Celnikier, D.. Successful first vaginal birth after cesarean section: a predictor of reduced risk for uterine rupture in subsequent deliveries. *Isr Med Assoc J*, July 2000; 2(7): 526-8.

Society of Obstetricians and Gynecologists of Canada (SOGC), *Guidelines for Vaginal Birth After Previous Cesarean Birth*. No. 155, February 2005.

Sternberg, C., McNiven, P. Midwifery Care for Women Planning Vaginal Birth After Cesarean Section: A survey of Ontario Midwifery Practices, *Canadian Journal of Midwifery Research and Practice*, Spring 2003; Vol. 2. No. 1, 10 – 14, Canadian Association of Midwives (CAM)

Wilson, K. Commentary: Vaginal Birth After Cesarean Section and the Issue of Home Birth, *Canadian Journal of Midwifery Research and Practice*, Spring 2003; Vol. 2. No. 1, 7 - 9, CAM

Zelop, C., Uterine Rupture During a Trial of Labour After Prior Cesarean Birth – Contemporary Insights. *European Obstetrics and Gynaecology*, 2012; 7(1): 32-4